

On the Congruency-Constrained Matroid Base ^{*}

Siyue Liu[†]

Chao Xu[‡]

Abstract

Consider a matroid where all elements are labeled with an element in \mathbb{Z} . We are interested in finding a base where the sum of the labels is congruent to $g \pmod{m}$. We show that this problem can be solved in $\tilde{O}(2^{4m}nr^{5/6})$ time for a matroid with n elements and rank r , when m is either the product of two primes or a prime power. The algorithm can be generalized to all moduli and, in fact, to all abelian groups if a classic additive combinatorics conjecture by Schrijver and Seymour holds true. We also discuss the optimization version of the problem.

1 Introduction

Recently, there has been a surge of work on congruency-constrained combinatorial optimization problems, such as submodular minimization [21], constraint satisfaction problems [5], and integer programming over totally unimodular matrices [1, 20]. In this paper, we consider congruency-constrained matroid base problems.

As a motivation, consider the exact matching problem, which asks for a perfect matching in a red-blue edge-colored bipartite graph with exactly k red edges [23]. A more general variant, where each edge has a weight of polynomial size and the goal is to find a perfect matching with an exact weight, has a randomized polynomial-time algorithm [18]. Exact matching is a special case of the exact matroid intersection problem, where the goal is to find a common base of a particular weight. Webb provided an algebraic formulation of the exact base problem that leads to an efficient solution when the matroid pair is Pfaffian and the weights are small [26]. For a single matroid, the exact base problem is defined as follows.

Problem 1 (Exact Matroid Base) *Given a matroid $M = (E, \mathcal{I})$, a natural number t , and a label function $\ell : E \rightarrow \mathbb{N}$. Find a base B in M such that $\sum_{x \in B} \ell(x) = t$, if one exists.*

The problem is NP-hard, as the classical subset sum problem can be reduced to it. However, the subset sum and its optimization variant, the knapsack problem, can be solved in pseudopolynomial time [3]. Thus, Papadimitriou and Yannakakis questioned whether the exact matroid base could be solved in pseudopolynomial time [23]. Barahona and Pulleyblank provided an affirmative answer for graphic matroids [2]. Subsequently, Camerini and Maffioli demonstrated that for matroids representable over the reals, there exists a randomized pseudopolynomial time algorithm for the exact matroid base problem [9]. Recently, Doron-Arad, Kulik, and Shachnai showed that pseudopolynomial time algorithms for arbitrary matroids cannot exist, demonstrated through carefully constructed paving matroids [12]. For one optimization variant, where the goal is to maximize the value of a base while adhering to an upper bound on the budget, some approximation results are known [7, 8].

^{*}We became aware after our submission that this problem was posed at 14th Emléktábla Workshop.

[†]siyueliu@andrew.cmu.edu, Carnegie Mellon University.

[‡]the.chao.xu@gmail.com, University of Electronic Science and Technology of China.

Recent progress has involved relaxing the exact constraint to a modular constraint to achieve interesting advancements. For instance, finding a perfect matching with an even number of red edges has been shown to be solvable in deterministic polynomial time [13]. Hence, we relax the exact constraint and study the following problem:

Problem 2 (m -Congruency-Constrained Matroid Base (CCMB(m))) *Let a matroid $M = (E, \mathcal{I})$, $g \in \{0, \dots, m-1\}$, and a label function $\ell : E \rightarrow \mathbb{Z}$. Find a base B in M such that $\sum_{x \in B} \ell(x) \equiv g \pmod{m}$, if any.*

As far as we know, except for results implied by the exact matroid base problem, little is known about this problem. An algorithm with a running time polynomial in m and the size of the matroid cannot exist, as it would imply the existence of a pseudopolynomial time algorithm for the exact matroid base problem. We aim for the next best thing: does there exist a fixed-parameter tractable (FPT) algorithm for CCMB(m), parameterized by m ?

Similar to [19], we consider finite abelian groups instead of solely congruency constraints. For a finite abelian group G , let $\ell : E \rightarrow G$ be a label function mapping each element of the matroid to an element of G . ℓ is called a G -labeling of E . A base with label sum $\sum_{x \in B} \ell(x) = g$ is called a g -base. Our goal can be summarized as determining the complexity of the following two problems regarding finding g -bases, parameterized by some finite abelian group G .

Problem 3 (Group-Constrained Matroid Base (GCMB(m))) *Given $g \in G$, a matroid $M = (E, \mathcal{I})$, and a label function $\ell : E \rightarrow G$, find a g -base, if one exists.*

Problem 4 (Group-Constrained Optimum Matroid Base (GCOMB(m))) *Given $g \in G$, a matroid $M = (E, \mathcal{I})$, a label function $\ell : E \rightarrow G$, and a weight function $w : E \rightarrow \mathbb{R}$, find a g -base of minimum weight, if one exists.*

The closest study related to matroid bases and group constraints comes from the additive combinatorics community, where theorems considering the number of different labels attainable by the bases have been discovered [11, 24].

Our Contribution. We show that GCMB(m) can be solved in $\tilde{O}(2^{4|G|} nr^{5/6})$ time for a group G , if either the size of G equals the product of two primes, or G is a cyclic group of size a power of a prime. In particular, this proves that CCMB(m) is in FPT parameterized by m if m is the product of two primes or a prime power.

These results are obtained through proximity results. For this, let \mathcal{B} be the set of bases of M . Let $\mathcal{B}(g)$ be the collection of all g -bases of M . For a weight function $w : E \rightarrow \mathbb{R}$, denote by $\mathcal{B}^* := \arg \min_{B \in \mathcal{B}} w(B)$ and $\mathcal{B}^*(g) := \arg \min_{B \in \mathcal{B}(g)} w(B)$ the sets of *optimum bases* and *optimum g -bases*, respectively. We introduce the following key concepts of this paper.

Definition A G -labeling ℓ of matroid M is **k -close** if for every $g \in G$ such that a g -base exists, every base has a g -base differing from it by at most k elements. If every G -labeling is k -close for every matroid, then G is **k -close**.

Definition A G -labeling ℓ of matroid M is **strongly k -close** if for every weight $w : E \rightarrow \mathbb{R}$ and $g \in G$ such that a g -base exists, every optimum base has an optimum g -base differing from it by at most k elements. If every G -labeling is strongly k -close for every matroid, then G is **strongly k -close**.

Note that being strongly k -close implies being k -close by setting $w \equiv 0$. We prove that G is $(|G|-1)$ -close if either the size of G equals the product of two primes, or G is a cyclic group of size a power

of a prime (Corollary 5.8). More generally, we show that every group is $(|G| - 1)$ -close assuming a conjecture by Schrijver and Seymour (Conjecture 5.5). This gives rise to the desired FPT algorithm for $\text{GCMB}(m)$ (Theorem 3.1). Lastly, we prove the strong k -closeness for strongly base orderable matroids (Theorem 6.1) and small groups, for some $k \leq |G|$ which only depends on G . These lead to FPT algorithms for $\text{GCOMB}(m)$ for those matroids and groups.

Overview. In Section 2, we define the problem and provide the necessary background in matroid and group theory. In Section 3, we discuss established algorithms and demonstrate the existence of an FPT algorithm for $\text{GCMB}(m)$ and $\text{GCOMB}(m)$, provided that G is k -close and strongly k -close, respectively, for some k which only depends on G . We deal with the k -closeness in Section 4 and 5. In Section 4, we consider what would constitute the minimum counterexample to the statement that G is k -close. We observe that the minimum counterexample would have to be a block matroid of rank $k + 1$. In Section 5, we establish the $(|G| - 1)$ -closeness properties for certain groups. This is achieved by connecting to an additive combinatorics theorem by Schrijver and Seymour. Finally, Section 6 discusses the optimization version and results on strong k -closeness.

Notation. We are given a matroid $M = (E, \mathcal{I})$, a group G , and a label function $\ell : E \rightarrow G$. For any $F \subseteq E$, denote by $\ell(F) := \sum_{e \in F} \ell(e)$. For a subset $H \subseteq G$, denote by $E(H) := \ell^{-1}(H) = \{e \in E \mid \ell(e) \in H\}$ all the elements with labels in H . If $H = \{g\}$ is a singleton, we shorten $E(\{g\})$ to $E(g)$. Denote by $\ell(M)$ the set $\{\ell(B) \mid B \text{ is a base of } M\}$. For $F \subseteq E$, denote the rank of F in M by $r_M(F)$, where M in the rank function $r_M(\cdot)$ can be omitted if the context is clear. Denote the rank of matroid M by $r(M) := r_M(E)$.

2 Preliminaries

Let $M = (E, \mathcal{I})$ be a matroid, where E is the ground set and \mathcal{I} is the family of independent sets. For a subset $F \subseteq E$, denote by $M \setminus F$ the *deletion* minor of M defined on the ground set $E \setminus F$ whose independent sets are those of M restricted to $E \setminus F$. If $F \subseteq E$ is an independent set of M , denote by M/F the *contraction* minor of M defined on the ground set $E \setminus F$. A subset $A \subseteq E \setminus F$ is independent in M/F if and only if $A \cup F$ is independent in M . Their rank functions satisfy the relation $r_{M/F}(A) = r_M(A \cup F) - r_M(F)$. We assume the matroid is loopless throughout, which means $\{e\} \in \mathcal{I}$ for any $e \in E$.

A matroid possesses the base exchange property: For any two bases A and B , there exists a series of elements $a_1, \dots, a_k \in A \setminus B$ and $b_1, \dots, b_k \in B \setminus A$ such that $A - \{a_1, \dots, a_j\} + \{b_1, \dots, b_j\}$ is also a base for each $j = \{1, \dots, k\}$, and $A - \{a_1, \dots, a_k\} + \{b_1, \dots, b_k\} = B$. The distance $d(A, B)$ between two bases A and B is the number of steps needed to exchange elements from A to B , which equals $|A \setminus B|$.

A matroid is *strongly base orderable* if for any two bases A and B , there is a bijection $f : A \rightarrow B$ such that for any $X \subseteq A$, we have $A - X + f(X)$ is a base. A function with the previous property can be taken to satisfy $f(a) = a$ for all $a \in A \cap B$. A matroid is a *block matroid* if the ground set is the union of two disjoint bases, and such disjoint bases are called *blocks*.

Given a group G and a normal subgroup $H \subseteq G$, denote by G/H the quotient group consisting of equivalence classes of the form $gH := \{g + h \mid h \in H\}$, which are called the *cosets* of H , where g_1 is equivalent to g_2 if and only if $g_1^{-1}g_2 \in H$. For a subset $F \subseteq G$, the *stabilizer* of F is defined by $\text{stab}(F) := \{g \in G \mid g + F = F\}$. It is easy to see that $\text{stab}(F)$ is a subgroup of G .

The Davenport constant [10, 22] of G , denoted by $D(G)$, is the minimum value such that every sequence of elements from G of length $D(G)$ contains a non-empty subsequence that sums to 0. In other words, the longest sequence without non-empty subsequence that sums to 0 has length $D(G) - 1$. It holds that $D(G) \leq |G|$, and equality holds when G is cyclic [25].

Our paper makes progress on the following two conjectures.¹

¹In an earlier preprint version of this paper, we conjectured that every finite abelian group is (strongly) $(D(G) - 1)$ -close, but this was recently disproved by [14].

Conjecture 2.1 (Feasibility) *Every finite abelian group G is $(|G| - 1)$ -close.*

Conjecture 2.2 (Optimization) *Every finite abelian group G is strongly $(|G| - 1)$ -close.*

Observe the conjectures are best possible when the group is cyclic. Indeed, the tight example is a block matroid of size $|G| - 1$, where the first block has all its element labeled 1 and the second has all its elements labeled 0. Then, the closest 0-base to the first block is the second block, which has distance $|G| - 1$ from it.

3 Algorithms for GCOMB

In this section, we show how k -closeness and strong k -closeness of a group G lead to FPT algorithms for GCMB(m) and GCOMB(m), respectively. In particular, Theorem 3.1 shows that if there is a function f such that G is $f(G)$ -close or strongly $f(G)$ -close, then for a fixed group G , GCMB(m) or GCOMB(m) is in FPT, respectively.

We assume that the matroid can be accessed through an independence oracle that takes a constant time per query. Each group operation also takes $O(1)$ time.

There is a folklore algorithm which shows for any fixed group G , GCOMB(m) is solvable in polynomial time by reducing it to a polynomial number of matroid intersections.

Let $M = (E, \mathcal{I})$ be a matroid of size n and rank r . Recall $E(g) = \ell^{-1}(g)$ for $g \in G$. For a base B of matroid M , the vector $a \in \mathbb{Z}_{\geq 0}^G$ such that $a_g = |B \cap E(g)|$ is called the *signature* of B . For a target label $h \in G$, every h -base B satisfies $h = \sum_{g \in G} a_g \cdot g$ for the signature a of B . Here $a \cdot g$ is defined to be $\sum_{i=1}^a g$.

For a fixed vector a with $\sum_g a_g = r$, $0 \leq a_g \leq |E(g)|$ and $\sum_g a_g \cdot g = h$, we can check if any base has signature a , in fact, find a minimum cost base with signature a . Indeed, one can obtain this through matroid intersection with a partition matroid, whose partition classes are $\{E(g) \mid g \in G\}$. It suffices to find an optimum base of matroid M such that its intersection with $E(g)$ has cardinality a_g for each $g \in G$. So the number of vectors we need to check is bounded above by $\binom{r+|G|-1}{|G|-1} \leq r^{|G|-1}$. Hence, we have to run the matroid intersection algorithm $r^{|G|-1}$ times in order to find the optimum g -base. Currently, the fastest algorithm for matroid intersection has running time $O(nr^{5/6} \log r)$ time assuming independence oracle takes constant time [4]. Hence the total running time is $\tilde{O}(nr^{O(|G|)})$ which is polynomial as $|G|$ is fixed. Although the algorithm is polynomial time for a fixed G , if we parameterize by G , then this algorithm is not FPT.

If G is (strongly) k -close for a constant k , then there is also a polynomial time algorithm: find an (optimum) base B , consider all possible bases B' such that $|B \setminus B'| \leq k$, and return an (optimum) base B' with the desired label. The running time is $O(r^k n^k)$ by enumerating all bases k -close to B .

We show that (strong) k -closeness gives rise to an FPT algorithm parametrized by G by combining it with the matroid intersection approach.

Theorem 3.1 *If G is k -close or strongly k -close, then there is a $\tilde{O}\left(\binom{k+|G|-1}{k}^2 nr^{5/6}\right)$ running time algorithm for GCMB(m) or GCOMB(m), respectively.*

Proof: The algorithm consists of two parts. First find any base B . By (strong) k -closeness, we know there is a feasible (optimum) g -base D that differs from B in at most k positions. Let a_D and a_B be the signature vectors of D and B , respectively. Let $a^+ := (a_D - a_B) \vee 0$, $a^- := (a_D - a_B) \wedge 0$ be vectors taking coordinate-wise maximum and minimum of $(a_D - a_B)$ with 0, respectively. Note that $a^+ \geq 0$ and $\sum_g a_g^+ \leq k$. Therefore, there are at most $\binom{k+|G|-1}{k}$ possible a^+ 's, and similarly at most $\binom{k+|G|-1}{k}$ possible a^- 's. Hence, we only need to run the matroid intersection algorithm for at most $\binom{k+|G|-1}{k}^2$ times to search for the (optimum) g -bases which are (strongly) k -close to B . \square

If we take $k = |G| - 1$ and use the fact that $\binom{2k}{k} \leq 2^{2k}$, we obtain the following corollary.

Corollary 3.2 *If G is $(|G| - 1)$ -close or strongly $(|G| - 1)$ -close, then there is an algorithm whose running time is $\tilde{O}(2^{4|G|} n r^{5/6})$ for GCMB(m) or GCOMB(m), respectively.*

4 Properties of minimum counterexamples to Conjecture 2.1 and Conjecture 2.2

In this section, we show that to prove Conjecture 2.1 or Conjecture 2.2, it suffices to prove it for a restricted type of matroids. Assume for the sake of contradiction that G is not (strongly) k -close, then there exists a counterexample of minimum size. This section shows that the minimum counterexamples are block matroids of size $(k + 1)$. Hence, it suffices to prove the conjectures for those matroids. We need a lemma by Brualdi [6].

Lemma 4.1 ([6]) *Let A and B be bases in a matroid, then there exists a bijection $f : A \setminus B \rightarrow B \setminus A$, such that $A - a + f(a)$ is a base.*

We give the following lemma.

Lemma 4.2 *Given a matroid $M = (E, \mathcal{I})$ and any weight function $w : E \rightarrow \mathbb{R}$, let A be an optimum base such that its weight is minimized. Then, for any $a \in A$, there exists some $b \in E \setminus A$ such that $A - a + b$ is an optimum base in $M' = M \setminus a$ with weight being the restriction of w to $E \setminus a$.*

Proof: Suppose A' is an optimum base in $M \setminus a$. By Lemma 4.1, there exists a bijection $f : A \setminus A' \rightarrow A' \setminus A$ such that $A - e + f(e)$ is a base for any $e \in A \setminus A'$. Since A is an optimum base, we have $w(A) \leq w(A - e + f(e))$, and thus $w(e) \leq w(f(e))$ for each $e \in A \setminus A'$. For any $a \in A$, let $b = f(a)$. Then, $w(A - a + b) = w(A) - w(a) + w(b) \leq w(A) - \sum_{e \in A \setminus A'} (w(e) - w(f(e))) = w(A')$. The inequality is because $a \in A \setminus A'$ and $w(e) \leq w(f(e))$ for any $e \in A \setminus A' \setminus a$. Since A' is an optimum base in $M \setminus a$, $A - a + b$ is also an optimum base in $M \setminus a$. \square

Next, we show that if G is not strongly k -close, then there has to be a G -labeling ℓ , a weight function w , an element $g \in G$, and a block matroid with blocks A, B , such that A is an optimum base, B is the closest optimum g -base to A , and $d(A, B) > k$. We call the pair of bases A, B a *witness*.

Theorem 4.3 *If a finite abelian group G is not strongly k -close, then there is a block matroid M of rank $k + 1$ with blocks A and B that form a witness.*

Proof: Let matroid $M = (E, \mathcal{I})$ be the smallest matroid, in terms of the number of elements, such that G is not strongly k -close for M . Let $w : E \rightarrow \mathbb{Z}$ be a weight function and $g \in G$. Recall that $\mathcal{B}^* := \arg \min_{B \in \mathcal{B}} w(B)$ and $\mathcal{B}^*(g) := \arg \min_{B \in \mathcal{B}(g)} w(B)$ are the sets of optimum bases and optimum g -bases, respectively. There exists $A \in \mathcal{B}^*$ and $B \in \mathcal{B}^*(g)$ such that B is an optimum g -base closest to A but $|A \setminus B| > k$.

First, we argue that $A \cap B = \emptyset$. If not, let $M' = (E', \mathcal{I}')$ be the matroid obtained by contracting $A \cap B$, and let $A' = A \setminus B$, $B' = B \setminus A$, $g' = g - \ell(A \cap B)$ and $w' = w|_{E'}$. Clearly, A' is a weighted minimum base of M' and B' is a weighted minimum g' -base of M' with weight w' . Also, $d(A', B') = |A' \setminus B'| = |A \setminus B| > k$, and B' is an optimum g' -base closest to A' . However, $|E'| = |E| - |A \cap B| < |E|$, contradicting to the minimality of $|E|$. Moreover, $A \cup B = E$. Otherwise, let M' be the matroid obtained by deleting $E \setminus (A \cup B)$ and A, B stays a witness in M' , contradicting to the minimality of $|E|$. Therefore, M is a block matroid which is a block matroid with blocks A and B .

Suppose $r(M) > k + 1$. Pick any $a \in A$. Let $M' = (E', \mathcal{I}') := M \setminus a$. By Lemma 4.2, there exists $b \in B$ such that $A' := A - a + b$ is a weighted minimum base in M' . Clearly, B is still a weighted minimum

g -base in M' that is closest to A' . Moreover, $d(A', B) = |A' \setminus B| = |A| - 1 = r(M) - 1 > k$. Therefore, G is not strongly k -close for M' , since A', B is a witness. Because $|E'| < |E|$, we obtain a contradiction. Therefore, $r(M) \leq k + 1$. Combining this with the fact that $r(M) = |A \setminus B| > k$, we get $r(M) = k + 1$. \square

As a corollary, if G is not k -close, then there has to be a G -labeling ℓ , an element $g \in G$ and a block matroid M with blocks A, B , such that A is a base, B is the closest g -base to A , and $d(A, B) > k$. Similarly, we call the pair of bases A, B a *witness* in the unweighted setting.

Corollary 4.4 *If a finite abelian group G is not k -close, then there is a block matroid M of rank $k + 1$ with blocks A and B that form a witness.*

Proof: The proof follows by setting the weight function to be 0 in the proof of Theorem 4.3. \square

Therefore, in order to show G is k -close or strongly k -close, we just have to show it is k -close or strongly k -close for all block matroids of rank $k + 1$, respectively.

5 k -closeness and isolation

We discuss progress towards conjecture Conjecture 2.1 in this section. We will prove in Theorem 5.6 that whenever a certain additive combinatorics conjecture is true, any finite abelian group G is $(|G| - 1)$ -close. First, we set up some notions we use later. We say a base B is *isolated* under label ℓ if it is the unique base with the label $\ell(B)$. A labeling is called *block isolating* if it isolates a base whose complement is also a base, i.e., it isolates a block. The next proposition shows that isolation and k -closeness are related concepts.

Proposition 5.1 *If no G -labeling of a rank $k + 1$ block matroid M is block isolating, then G is k -close for M .*

Proof: Suppose not. By Corollary 4.4, there exists a G -labeling ℓ , $g \in G$ and witness A, B that are bases of M such that B is the closest g -base to A . Then, B is the unique base that has labeling g . Indeed, if there is any other base $B' \neq B$ with $\ell(B') = g$, then $d(A, B') < k + 1 = d(A, B)$, since the block structure guarantees that B is the only farthest base from A , a contradiction. \square

Equipped with Proposition 5.1, our goal becomes showing that block isolating labelings cannot exist.

5.1 Congruency-constrained base with prime modulus

We will start by proving that Conjecture 2.1 is true for any cyclic group of prime order, or equivalently congruency-constraints modulo primes. This is a special case of results for general groups which will be introduced in the next subsection. But the proof for cyclic groups of prime order is much simpler followed by a counting argument. For the sake of helping readers gain intuition, we present it as well. The main tool we are going to use is the following additive combinatorics lemma by Schrijver and Seymour.

Lemma 5.2 (Schrijver-Seymour [24]) *Let $M = (E, \mathcal{I})$ be a matroid with rank function r and let $\ell : E \rightarrow \mathbb{Z}_p$ for some prime number p . Let $\ell(M) := \{\ell(B) \mid B \text{ is a base of } M\}$. Then, $|\ell(M)| \geq \min\{p, \sum_{g \in \mathbb{Z}_p} r(E(g)) - r(M) + 1\}$.*

The following lemma states an exchange property for matroids which is most likely routine. We also give its proof here.

Lemma 5.3 *Given a matroid $M = (E, \mathcal{I})$, let A be a base, $A_1 \subseteq A$, and B_1 be an independent set such that $A \cap B_1 = \emptyset$. If $|A_1| + |B_1| - r(A_1 \cup B_1) \geq t$, then there exist some $A_2 \subseteq A_1$ and $B_2 \subseteq B_1$ with $|A_2| = |B_2| = t$, such that $A - A_2 + B_2$ is a base.*

Proof: By submodularity of matroid rank functions,

$$r((A \setminus A_1) \cup B_1) + r(A_1 \cup B_1) \geq r(A \cup B_1) + r(B_1) = r(M) + |B_1|.$$

By assumption, $|A_1| + |B_1| - r(A_1 \cup B_1) \geq t$. Combining these, we have

$$r((A \setminus A_1) \cup B_1) \geq r(M) - |A_1| + t = |A \setminus A_1| + t.$$

Using the fact that $A \setminus A_1$ is independent, we deduce there exists $B_2 \subseteq B_1$ with $|B_2| = t$, such that $(A \setminus A_1) \cup B_2$ is independent. Moreover, since A is a base, by adding elements from A_1 , $(A \setminus A_1) \cup B_2$ can be extended to a base which has the form $A - A_2 + B_2$ for some $A_2 \subseteq A_1$ with $|A_2| = |B_2| = t$. \square

Theorem 5.4 *For any prime p , \mathbb{Z}_p is $(p-1)$ -close.*

Proof: Suppose not. Then, by Proposition 5.1, there exists a block matroid $M = (E, \mathcal{I})$ with $E = A \cup B$, where A and B are two disjoint bases with $|A| = |B| = p$, such that A is isolated under some \mathbb{Z}_p -labeling ℓ .

We claim that for any $g \in G$, $r(E(g)) = |E(g) \cap A| + |E(g) \cap B| = |E(g)|$.

Otherwise, letting $A_1 = E(g) \cap A$, $B_1 = E(g) \cap B$ and $t = 1$ in Lemma 5.3, we can find $a \in A_1$ and $b \in B_1$ such that $A - a + b$ is a base. Since $\ell(a) = \ell(b) = g$, $\ell(A - a + b) = \ell(A)$, contradicting to the fact that A is isolated.

Take any element $e \in A$ and consider the matroid $M' = M \setminus e$ with ground set $E' = E \setminus e$. Note that $r(M') = r(M)$ because B stays a base of M' . Applying Lemma 5.2 to M' ,

$$\begin{aligned} |\ell(M')| &\geq \min \left\{ p, \sum_{g \in \mathbb{Z}_p} r(E'(g)) - r(M') + 1 \right\} \\ &= \min \left\{ p, \sum_{g \in \mathbb{Z}_p, g \neq \ell(e)} r(E(g)) + r(E(\ell(e)) \setminus \{e\}) - r(M) + 1 \right\} \\ &= \min \left\{ p, \sum_{g \in \mathbb{Z}_p, g \neq \ell(e)} |E(g)| + (|E(\ell(e))| - 1) - r(M) + 1 \right\} \\ &= \min \left\{ p, \sum_{g \in \mathbb{Z}_p} |E(g)| - p \right\} = \min \{ p, 2p - p \} = p. \end{aligned}$$

Thus, there exists a base D of M' such that $\ell(D) = \ell(A)$. By definition, D is also a base of M . Since $e \in A$ but $e \notin D$, D is distinct from A , contradicting to the fact that A is isolated under label ℓ . \square

5.2 General group constraints

To extend beyond prime modulus, we introduce the following conjecture of Schrijver and Seymour which is an extension of Lemma 5.2 to any abelian group G .

Conjecture 5.5 (Schrijver-Seymour [24], see also [11]) *Let $M = (E, \mathcal{I})$ be a matroid with rank function r and let $\ell : E \rightarrow G$ for some finite abelian group G . Let $\ell(M) := \{\ell(B) \mid B \text{ is a base of } M\}$. Let $H = \text{stab}(\ell(M))$ be the stabilizer of $\ell(M)$. Then, $|\ell(M)| \geq |H| \cdot \min \left\{ \sum_{Q \in G/H} r(E(Q)) - r(M) + 1, |G|/|H| \right\}$.*

For a prime p and a cyclic group $G = \mathbb{Z}_p$, if $\ell(M) \neq \mathbb{Z}_p$, then it is easy to see $\text{stab}(\ell(M)) = \{0\}$. Then, the inequality in Conjecture 5.5 reduces to $|\ell(M)| \geq \min \left\{ \sum_{g \in G} r(E(g)) - r(M) + 1, |G| \right\}$, which is precisely the form in Lemma 5.2. Otherwise, $\text{stab}(\ell(M)) = \mathbb{Z}_p$ and the inequality trivially holds.

Theorem 5.6 *If Conjecture 5.5 is true for a finite abelian group G and all of its subgroups, then G is $(|G| - 1)$ -close.*

Proof: Let $n := |G|$. We proceed by induction on n . The theorem trivially holds when $n = 1$. Suppose the theorem does not hold for some $|G| = n$. Then by Proposition 5.1, there exists a block matroid $M = (E, \mathcal{I})$, $E = A \cup B$, where A and B are two disjoint bases of M with $|A| = |B| = n$ such that A is isolated under some labeling ℓ .

Take any $e \in A$, let $M' = (E', \mathcal{I}') := M \setminus e$. Let H be the stabilizer of $\ell(M')$. Since G is abelian, H is a normal subgroup of G . Denote by $gH := \{g + h \mid h \in H\}$ the coset of H with representative g . First, observe that if $g \in \ell(M')$, then $gH \subseteq \ell(M')$. This is because, by definition of H , for any $h \in H$, $g + h \in \ell(M')$. Thus $gH \subseteq \ell(M')$. Therefore, $\ell(M') = \bigcup_{g \in R} gH$, where R is a collection of representatives of cosets of H . It follows that $|\ell(M')| = |R| \cdot |H|$. Let $E'(gH) = E(gH) \setminus \{e\}$ for any $g \in G$. Since A is isolated under ℓ and A is not a base of M' , we know $\ell(A) \notin \ell(M')$. Thus, $|\ell(M')| < |G|$. This implies $|H| < |G|$, since $|G| > |\ell(M')| = |R| \cdot |H| \geq |H|$ as $R \neq \emptyset$. It also follows from $|\ell(M')| < |G|$ that $|\ell(M')| \geq |H|$ ($\sum_{Q \in G/H} r(E'(Q)) - r(M') + 1$) in Conjecture 5.5, and thus $|R| \geq \sum_{Q \in G/H} r(E'(Q)) - r(M') + 1 = \sum_{Q \in G/H} r(E'(Q)) - n + 1$. Then,

$$\begin{aligned} & \sum_{Q \in G/H} (|A \cap E'(Q)| + |B \cap E'(Q)| - r(E'(Q))) \\ &= (2n - 1) - \sum_{Q \in G/H} r(E'(Q)) \\ &\geq (2n - 1) - (n + |R| - 1) \\ &= n - |R|. \end{aligned}$$

Therefore,

$$\begin{aligned} & \max_{Q \in G/H} (|A \cap E'(Q)| + |B \cap E'(Q)| - r(E'(Q))) \\ &\geq \frac{n - |R|}{|G/H|} \\ &> \frac{n - |G/H|}{|G/H|} \\ &= |H| - 1, \end{aligned}$$

where the strict inequality follows from $|\ell(M')| < |G|$. Suppose the maximum is attained at $Q_0 = g_0H$. Then, we have

$$|A \cap E'(Q_0)| + |B \cap E'(Q_0)| - r(E'(Q_0)) \geq |H|,$$

since $|A \cap E'(Q_0)| + |B \cap E'(Q_0)| - r(E'(Q_0))$ is an integer. Let $A_1 := A \cap E'(Q_0)$, $B_1 := B \cap E'(Q_0)$. By Lemma 5.3, there exists a base $A - A_2 + B_2$ such that $A_2 \subseteq A_1$, $B_2 \subseteq B_1$ and $|A_2| = |B_2| = |H|$.

Let $M'' = (E'', \mathcal{I}'')$ be the matroid obtained by contracting $A \setminus A_2$ and deleting $B \setminus B_2$, i.e. $M'' = M / (A \setminus A_2) \setminus (B \setminus B_2)$. Note that M'' is a rank $|H|$ block matroid with blocks A_2 and B_2 . Consider an H -labeling on $E'' = A_2 \cup B_2$, $\ell'' : E'' \rightarrow H$, such that $\ell''(e) = \ell(e) - g_0 \in H, \forall e \in E''$. Note that $|H| < |G|$. By the induction hypothesis and the assumption that Conjecture 5.5 is true for the subgroup H of G , we know that there is no isolating H -labeling of M'' . Thus, there must be another base $D_2 \neq A_2$ of M'' , such that $\ell''(D_2) = \ell''(A_2)$. Thus $\ell(D_2) = \ell''(D_2) + |H| \cdot g_0 = \ell''(A_2) + |H| \cdot g_0 = \ell(A_2)$. By the definition of contraction, $D := D_2 \cup (A \setminus A_2)$ is a base of matroid M which is distinct from A . And $\ell(D) = \ell(D_2) + \ell(A \setminus A_2) = \ell(A_2) + \ell(A \setminus A_2) = \ell(A)$, contradicting to the fact that A is isolated under label ℓ . \square

The following lemma is the current status showing on which group the Schrijver-Seymour conjecture is true.

Lemma 5.7 ([11]) *Conjecture 5.5 is true for G if*

1. $|G| = pq$ for primes p and q , or
2. $G = \mathbb{Z}_{p^n}$ for a prime p and a positive integer n .

Corollary 5.8 G is $(|G| - 1)$ -close if $|G| = pq$, or $G = \mathbb{Z}_{p^n}$ for any p, q primes and n a positive integer.

6 Strong k -closeness

This section discusses our results for strong k -closeness. Unfortunately, we have little understanding of strong k -closeness. Hence, we focus on restricted class of matroids and small groups.

For the matroids and groups concerned within this section, we obtain a tighter bounds than Conjecture 2.2 implies. Instead of $|G| - 1$ in the conjecture, we have $D(G) - 1$.

6.1 Strongly base orderable matroids

We first study strongly base orderable matroids.

Theorem 6.1 *Every G -labeling is strongly $(D(G) - 1)$ -close for strongly base orderable matroids.*

Proof: Suppose not. By noting that strongly base orderability is closed under taking minors [15], it is not hard to see from Theorem 4.3 that there exists a counterexample which is a rank $D(G)$ strongly base orderable block matroid M with blocks A and B as a witness. Let ℓ be a G -labeling of M , w be a weight function such that A is an optimum base and B is the optimum g -base closest from A .

Let $f : B \rightarrow A$ be the bijection obtained from strongly base orderability. Define $\ell'(X) = \ell(f(X)) - \ell(X)$ for all $X \subseteq B$. Since $|B| = D(G)$, by the definition of Davenport's constant, there is $B_1 \subseteq B$ such that $\ell'(B_1) = 0$. Let $A_1 = f(B_1)$. This implies $\ell(A_1) = \ell(B_1)$ and thus the base $B - B_1 + A_1$ has the same label as B , i.e. $B - B_1 + A_1$ is also a g -base.

Now, consider $A_2 = A - A_1$ and $B_2 = B - B_1$. One has $f(B_2) = A_2$ and therefore $B - B_2 + A_2 = A - A_1 + B_1$ is also a base. Since A is an optimum base, $w(A - A_1 + B_1) \geq w(A)$, which means $w(A_1) \leq w(B_1)$. Therefore, $w(B - B_1 + A_1) \leq w(B)$. This means $B - B_1 + A_1$ is also an optimum g -base. But $A_1 \subsetneq A$ since $\ell(A) \neq \ell(B)$, which means $B - B_1 + A_1$ is an optimum g -base closer to A , a contradiction. \square

In fact, we can define a weaker property. Two bases A, B are k -replaceable if there exists a bijection $f : B \setminus A \rightarrow A \setminus B$ such that for any subset $B' \subseteq B \setminus A$ of size at most k , $B - B' + f(B')$ is a base. We say a matroid is k -replaceable if every pair of bases are k -replaceable. We have that Theorem 6.1 also holds for $(D(G) - 1)$ -replaceable matroids.

6.2 Small groups

Next, we consider small groups. Theorem 4.3 shows one only needs to check block matroids of rank $D(G)$ to know if G is strongly $(D(G) - 1)$ -close.

For $G = \mathbb{Z}_2$ which has $D(G) = 2$, observe that all size 4 matroids are strongly base orderable, hence \mathbb{Z}_2 is 1-close. For $G \in \{\mathbb{Z}_3, \mathbb{Z}_2 \times \mathbb{Z}_2\}$, we need to check all rank 3 block matroids. Since all but $M(K_4)$, the graphic matroid on K_4 , are strongly base orderable [17], we only need to test a single matroid. However, testing strongly 2-closeness of a single matroid is still computational intensive.

Instead, we introduce the notion of strong block isolation. A G -labeling ℓ *strongly isolates* a block B , if it is the unique block with label $\ell(B)$. A G -labeling is strong block isolating, if it strongly isolates a block.

Proposition 6.2 *If no G -labeling of a rank $k + 1$ block matroid M is strong block isolating, then G is strongly k -close for M .*

Proof: Suppose not. By Theorem 4.3, there exists a G -labeling ℓ , $g \in G$ and witness A, B that are bases of M such that B is the closest optimum g -base to A . Fix such labeling ℓ . Since ℓ is not strong block isolating, there is a non-empty $A_1 \subseteq A$ and $B_1 \subseteq B$ such that $A - A_1 + B_1$ is a block with the same label as A . Let $A_2 = A \setminus A_1$, $B_2 = B \setminus B_1$. Then the complement of $A - A_1 + B_1$ is the base $A - A_2 + B_2$, and has the same label as B . Since $w(B) = w(A) - w(A_1) + w(B_1) - w(A_2) + w(B_2) \geq w(A) - w(A_2) + w(B_2) = w(A - A_2 + B_2)$, $A - A_2 + B_2$ is a closer optimum g -base to A , a contradiction. \square

Hence, for each G in hand, we show no G -labeling of a rank $D(G)$ block matroid is strong block isolating, which implies G is strongly $(D(G) - 1)$ -close. We tested $M(K_4)$, and shows $\mathbb{Z}_3, \mathbb{Z}_2 \times \mathbb{Z}_2$ are strongly 2-close. Further, we test for each \mathbb{Z}_4 -labeling of a rank 4 block matroid. There are 940 non-isomorphic rank 4 matroids [16], and for each one, we test $4^8 = 65536$ different labeling. None of them is strongly block isolating. Therefore, we get the following.

Proposition 6.3 G is strongly $(D(G) - 1)$ -close if $|G| \leq 4$.

Acknowledgements

We thank the reviewers for their valuable feedback and András Imolay for noticing and correcting an error in Theorem 5.6.

References

- [1] Stephan Artmann, Robert Weismantel, and Rico Zenklusen. A strongly polynomial algorithm for bimodular integer linear programming. In *Proceedings of the 49th Annual ACM SIGACT Symposium on Theory of Computing*, STOC 2017, page 1206–1219, New York, NY, USA, 2017. Association for Computing Machinery.
- [2] Francisco Barahona and William R Pulleyblank. Exact arborescences, matchings and cycles. 16(2):91–99.
- [3] Richard Bellman. Notes on the theory of dynamic programming iv - maximization over discrete sets. *Naval Research Logistics Quarterly*, 3(1-2):67–70, 1956.
- [4] Joakim Blikstad. Breaking $O(nr)$ for Matroid Intersection. In Nikhil Bansal, Emanuela Merelli, and James Worrell, editors, *48th International Colloquium on Automata, Languages, and Programming (ICALP 2021)*, volume 198 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 31:1–31:17, Dagstuhl, Germany, 2021. Schloss Dagstuhl – Leibniz-Zentrum für Informatik.
- [5] Joshua Brakensiek, Sivakanth Gopi, and Venkatesan Guruswami. Constraint Satisfaction Problems with Global Modular Constraints: Algorithms and Hardness via Polynomial Representations. *SIAM Journal on Computing*, 51(3):577–626, June 2022.
- [6] Richard A. Brualdi. Comments on bases in dependence structures. *Bulletin of the Australian Mathematical Society*, 1(2):161–167, 1969.
- [7] P M. Camerini, F Maffioli, and C. Vercellis. Multi-constrained matroidal knapsack problems. *Mathematical Programming*, 45(1):211–231, August 1989.
- [8] Paolo M. Camerini and Carlo Vercellis. The matroidal knapsack: A class of (often) well-solvable problems. *Operations Research Letters*, 3(3):157–162, 1984.
- [9] PM Camerini, G Galbiati, and F Maffioli. Random pseudo-polynomial algorithms for exact matroid problems. 13(2):258–273.

- [10] H Davenport. Midwestern conference on group theory and number theory. *Ohio State University*, 1966.
- [11] Matt DeVos, Luis Goddyn, and Bojan Mohar. A generalization of kneser’s addition theorem. *Advances in Mathematics*, 220(5):1531–1548, 2009.
- [12] Ilan Doron-Arad, Ariel Kulik, and Hadas Shachnai. Tight lower bounds for weighted matroid problems, 2023.
- [13] Nicolas El Maalouly, Raphael Steiner, and Lasse Wulf. Exact Matching: Correct Parity and FPT Parameterized by Independence Number. In Satoru Iwata and Naonori Kakimura, editors, *34th International Symposium on Algorithms and Computation (ISAAC 2023)*, volume 283 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 28:1–28:18, Dagstuhl, Germany, 2023. Schloss Dagstuhl – Leibniz-Zentrum für Informatik.
- [14] Florian Hörsch, András Imolay, Ryuhei Mizutani, Taihei Oki, and Tamás Schwarcz. Problems on group-labeled matroid bases, 2024.
- [15] Aubrey W Ingleton. Transversal matroids and related structures. *Higher Combinatorics: Proceedings of the NATO Advanced Study Institute held in Berlin (West Germany), September 1–10, 1976*, pages 117–131, 1977.
- [16] Yoshitake Matsumoto, Sonoko Moriyama, Hiroshi Imai, and David Bremner. Matroid enumeration for incidence geometry. 47(1):17–43.
- [17] Dillon Mayhew and Gordon F Royle. Matroids with nine elements. *Journal of Combinatorial Theory, Series B*, 98(2):415–431, 2008.
- [18] Ketan Mulmuley, Umesh V Vazirani, and Vijay V Vazirani. Matching is as easy as matrix inversion. In *Proceedings of the Nineteenth Annual ACM Symposium on Theory of Computing, STOC ’87*, page 345–354, New York, NY, USA, 1987. Association for Computing Machinery.
- [19] Martin Nägele, Christian Nöbel, Richard Santiago, and Rico Zenklusen. Advances on strictly δ -modular ips. In *Proceedings of the 24th Conference on Integer Programming and Combinatorial Optimization (IPCO ’23)*, pages 393–407, 2023.
- [20] Martin Nägele, Richard Santiago, and Rico Zenklusen. *Congruency-Constrained TU Problems Beyond the Bimodular Case*, pages 2743–2790.
- [21] Martin Nägele, Benny Sudakov, and Rico Zenklusen. Submodular Minimization Under Congruency Constraints. *Combinatorica*, 39(6):1351–1386, December 2019.
- [22] John E Olson. A combinatorial problem on finite abelian groups, i. *Journal of number theory*, 1(1):8–10, 1969.
- [23] Christos H. Papadimitriou and Mihalis Yannakakis. The complexity of restricted spanning tree problems. *J. ACM*, 29(2):285–309, apr 1982.
- [24] Alexander Schrijver and Paul D Seymour. Spanning trees of different weights. In *Polyhedral combinatorics*, page 281, 1990.
- [25] P van Emde Boas and D. Kruyswijk. *A Combinatorial Problem on Finite Abelian Groups III*. Mathematisch Centrum, Amsterdam. Afdeling Zuivers Wiskunde. [Publications]. Stichting Mathematisch Centrum, 1969.

[26] Webb, Kerri. *Counting Bases*. PhD thesis, 2004.